

Supplement

Table S1. Comparison of Patients with or without Legal Procedures

Variables [†]	Without legal procedure (n=464)	With legal procedure (n=57)
	n (%)	n (%)
Length of stay (days), M/median±SD	23/13±40	78/63±61
Married	64 (13.8)	6 (10.5)
Race		
White	134 (28.9)	14 (24.6)
Black	105 (22.6)	18 (31.6)
Hispanic	147 (31.7)	15 (26.3)
Asian	66 (14.2)	10 (17.5)
Employed	88 (19.0)	6 (10.5)
Sleep disturbance	291 (62.7)	20 (35.1)
Medical comorbidities	279 (60.1)	31 (54.4)
History of alcohol use	191 (41.2)	10 (17.5)
History of drug use	193 (41.6)	17 (29.8)
History of psychiatric hospitalization	364 (78.5)	49 (86.0)
Antipsychotic medication	383 (82.5)	53 (93.0)
Antidepressants	164 (35.3)	3 (5.3)
Psychiatric disorder		
Schizophrenia	105 (22.6)	29 (50.9)
Schizoaffective disorder	79 (17.0)	17 (29.8)
Major depressive disorder	86 (18.5)	0 (0.0)
Bipolar disorder	75 (16.2)	4 (7.0)
Discharge Dispositions		
Home	325 (70.0)	25 (43.9)
Shelter	70 (15.1)	12 (21.1)
Adult home/group home	47 (10.1)	10 (17.5)
Long term care facilities/rehab program	7 (1.5)	2 (3.5)
Transfer to state psychiatric hospitals	5 (1.1)	6 (10.5)
Supporting house for intellectual deficits	2 (0.4)	1 (1.8)
Drug/alcohol rehab for substance abuse	7 (1.5)	0 (0.0)
Readmission within 30 days	26 (5.6)	4 (7.0)

[†] Variables were selected from Table 1 with clinician's judgement or support from published literature. This analysis was conducted exploratorily without formal comparison/testing (i.e., no *p*-value computed).